

ACETAMINOPHEN‡ (Tylenol, Tempra, etc.)					
WEIGHT		INFANT'S ORAL SUSPENSION 160 mg/5ml	CHILDREN'S LIQUID 160 mg/5ml	CHEWABLE 80 mg Tabs	CAPSULE OR CHEWABLE 160 mg Tabs
6-11 lbs	3-5 kg	1.25ml*	¼ tsp	***	***
12-17 lbs	6-8 kg	2.5ml*	½ tsp	***	***
18-23 lbs	9-10 kg	3.75ml	¾ tsp	***	***
24-35 lbs	11-16 kg	5ml	1 tsp	2 tab	1 tab
36-47 lbs	17-21 kg	----	1 ½ tsp	3 tab	1 ½ tab
48-59 lbs	22-27 kg	----	2 tsp	4 tab	2 tab
60-71 lbs	28-32 kg	----	2 ½ tsp	5 tab	2 ½ tab
72-95 lbs	33-43 kg	----	3 tsp	6 tab	3 tab
96 lbs +	44 kg +	----	----	----	4 tab

‡ Doses may be administered 4 or 5 times daily or as directed by your doctor.

Do not exceed 5 doses in 24 hours.

Consult your physician if fever persists for more than 3 days or if pain continues for more than 5 days. In case of accidental overdosage, consult a physician or poison control center immediately.

* Do not use in infants under 2 months of age unless directed by a physician.

*** Do not use this form in this weight range unless directed by a physician.

IBUPROFEN‡ (Motrin, Advil etc.)					
Recommended for 6 months & above		INFANT DROPS 50 mg/1.25ml	SUSPENSION 100 mg/5 ml	CHEWABLE 50 mg	CHEWABLE OR CAPLETS 100 mg Tabs
WEIGHT					
(pounds)	(kilogram)				
12-17 lbs	6-8 kg	1.25ml	½ tsp	***	***
18-23 lbs	9-10 kg	1.875ml	¾ tsp	***	***
24-35 lbs	11-16 kg	1.875 + 0.625 ml	1 tsp	2 tab	1 tab
36-47 lbs	17-21 kg	----	1 ½ tsp	3 tab	1 ½ tab
48-59 lbs	22-27 kg	----	2 tsp	4 tab	2 tab
60-71 lbs	28-32 kg	----	2 ½ tsp	5 tab	2 ½ tab
72-95 lbs	33-43 kg	----	3 tsp	6 tab	3 tab
96 lbs +	44 kg +	----	4 tsp	8 tab	4 tab

‡ Doses may be administered with minimum intervals of 6 hours.

Consult your physician if fever persists for more than 3 days or if pain continues for more than 5 days. In case of accidental overdosage, consult a physician or poison control center immediately.

*** Do not use this form in this weight range unless directed by a physician.